

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b>  <b>A61K 9/16</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/24109</b>  <b>(43) International Publication Date:</b> 9 December 1993 (09.12.93)
<b>(21) International Application Number:</b> PCT/US93/05423 <b>(22) International Filing Date:</b> 4 June 1993 (04.06.93)  <b>(30) Priority data:</b> 07/893,288                      4 June 1992 (04.06.92)                      US  <b>(71) Applicant (for all designated States except US):</b> SMITH-KLINE BEECHAM CORPORATION [US/US]; Corporate Patents - U.S., UW 2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> BHARDWAJ, Sanjay [IN/US]; 44 Center Grove Road, L-15, Gateways, Randolph, NJ 07869 (US). HAYWARD, Marshall [US/GB]; St. George's Avenue, Weybridge, Surrey KT13 0DE (GB).		<b>(74) Agents:</b> SUTER, Stuart, R. et al.; SmithKline Beecham Corporation, Corporate Patents - U.S., UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).  <b>(81) Designated States:</b> CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> PALATABLE PHARMACEUTICAL COMPOSITIONS  <b>(57) Abstract</b>  <p>A pharmaceutical granular composition and method for taste masking bitter, unpleasant tasting drugs comprising a drug core and as a taste masking agent methacrylate ester copolymers. The method comprises coating the drug cores with separate layers of aqueous dispersions of the copolymers. Additionally, the coating composition may contain plasticizers and conventional excipients. The granules of the present invention can be used in the preparation of chewable tablets which have good palatability and bioavailability. Preferable copolymers are poly(ethylacrylate, methylmethacrylate) to which quaternary ammonium groups have been introduced to modify the permeability of the ester. The coating system of this invention releases the drug by diffusion and is influenced by drug solubility and media pH.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

10

## PALATABLE PHARMACEUTICAL COMPOSITIONS

15        This invention relates to a composition and method for taste masking bitter, unpleasant tasting medicaments and provides for an immediate release of the medicament after ingestion. More specifically, the invention relates to medicament cores coated with methacrylate ester copolymers which masks the bitter and unpleasant taste of the medicament. A method of preparing dosage units such as chewable tablets, 20        employing the coated medicament is also described.

      In a preferred embodiment of this invention, the composition comprises a chewable tablet containing cimetidine coated with separate layers of methacrylate ester copolymers. The tablets do not exhibit the bitter taste of cimetidine and provides for an immediate release of the cimetidine.

25

### Background of the Invention

      Cimetidine is a histamine H<sub>2</sub>-antagonist which has been shown to be useful in the treatment of duodenal, gastric, recurrent and stomal ulceration. It is also employed 30        in the management of patients who are at high risk from hemorrhage of the upper gastrointestinal tract.

      Cimetidine is known to be one of many medicaments to have a pronounced bitter taste. This is not usually a problem when the dosage form employed is a capsule or a tablet designed to be swallowed, thereafter to disintegrate upon reaching the 35        stomach. However, such dosage forms can be impractical when it is desired to administer a large amount of active ingredient, or to co-administer a relatively bulky

second active ingredient such as an antacid or alginate. Moreover, many individuals have difficulty in swallowing a solid dosage form

A conventional approach to administering relatively large amounts of active ingredient in a solid dosage form is by means of a chewable tablet, i.e., a tablet which  
5 disintegrates in the mouth upon being chewed. Such a tablet also circumvents the problem of a solid dosage being difficult to swallow.

It will be appreciated that a major requirement of such a dosage form is that it must be palatable, since an unpalatable formulation increases the risk of a patient neglecting to take the tablet. Such non-compliance with the dosing regimen will in turn  
10 delay or prevent the patient's recovery from the condition under treatment.

A further requirement of such a composition is that once the disintegrated tablet reaches the stomach, the individual particles should release the active ingredient rapidly and completely in order to ensure that substantially all of the active ingredient is absorbed; that is to say the formulation should be readily bioavailable.

15 In the case of cimetidine, because of its bitterness and solubility characteristics, the provision of such a dosage form represents a considerable problem.

Several solutions to the problem of taste masking pharmaceutical compositions have been attempted. For example, cimetidine granules have been coated with various compositions such as ethylcellulose and polyvinyl and acrylic polymers. One such  
20 proposal is disclosed in U.S. Patent No. 4,800,087 wherein a polymer mixture coating is employed. The mixture comprises a high temperature film forming copolymer of polymethacrylic acid ester and acrylic acid ester and a low temperature film forming copolymers consisting of methacrylic acid ester and styrene acrylate. U.S. patent 4,892,740 discloses pharmaceutical preparations having improved flavouring  
25 characteristics obtained by the drug being coated by a polymeric substance which is soluble in gastric juice.

### DESCRIPTION OF THE INVENTION

30 In accordance with this invention it has been unexpectedly discovered that when bitter, unpleasant tasting medicaments are coated with separate layers of specific methacrylate ester copolymers a core results which is palatable and has good dissolution characteristics in the stomach. The coated core results in a taste masked medicament without reducing its bioavailability.

The core particles which are to be coated can be composed of pure granular drug material or drug granules prepared in the conventional manner employing appropriate binding agents. The particle size of the core material is from about 180 to 420 microns. This range is preferable because larger particles tend to be ruptured during chewing and smaller particles present problems in the coating process. The particles should possess adequate hardness and friability characteristics to withstand attritive forces during the coating operation.

Exemplary of bitter, unpleasant tasting drugs applicable to the taste-masking compositive of this invention are histamine H<sub>2</sub>-antagonists, such as, for example, cimetidine, ranitidine, famotidine, nizatidine, etinidine, lupitidine, nifentidine, niperotidine, roxatidine, sulfotidine, tuvatidine and zaltidine; antibiotics, such as penicillin, ampicillin and erythromycin, acetaminophen; caffeine, dextromethorphan, diphenhydramine, theophylline, spironolactone and chlorpheniramine. The above drugs are not limiting but merely exemplary of unpleasant tasting drugs that may be employed in this invention.

In a preferred embodiment the taste masking compositions of this invention will contain cimetidine or caffeine.

When, cimetidine is employed the composition will contain from about 75 mg to 800 mg. Preferably it will containing from about 100 mg. to 400 mg. of cimetidine.

The coating material is selected from the wide range of copolymers available under the tradename of Eudragits. These copolymers are manufactured and marketed by Röhm Pharma of Darmstadt, Germany. Preferably, the coating composition is comprised of aqueous dispersions of ethyl acrylate (EA) and methyl methacrylate (MMA). A particularly preferred coating composition of this invention is an aqueous dispersion of poly(EA-MMA) to which hydrophilic quaternary ammonium groups have been introduced to modify the permeability of the ester. Copolymerization of the above poly(EA-MMA)ester with different ratios of trimethylammonioethyl methacrylate chloride (TAMCl) results in methacrylic ester copolymers of either high or low permeability. For example, a coating film of high permeability results when poly (EA-MMA-TAMCl) are present in a 1:2:0.2 ratio. This copolymer is available under the tradename of Eudragit RL30D. A coating film of low permeability results when poly (EA-MMA-TAMCl) are present in a ratio of 1:2:0.1. This product is available under the tradename of Eudragit RS30D. Both Eudragit RL30D and RS30D are sold as 30% aqueous dispersions.

The polymeric aqueous dispersions may also contain additives such as, plasticizers, pigments, talc and the like. Plasticizers are employed to assist in the film forming characteristics of the polymeric coating and also to provide greater integrity and elasticity to the films coat. Exemplary of plasticizers that may be employed in the coatings of this invention are triethyl citrate, triacetin, tributyl citrate, acetyl triethyl citrate, acetyl tributyl citrate, dibutyl phthalate, dibutyl sebacate, vinyl pyrrolidone and propylene glycol. The amount of plasticizer present in the aqueous dispersion may be from 5% to about 30%. Preferably, the plasticizer will be present in an amount of from about 15% to about 25% by weight of the dry polymers.

Exemplary of pigments employed in the polymeric coating solutions of the invention may be color pigments such as FD&C or D&C approved lakes and dyes, iron oxide and titanium dioxide. The amount of pigment present in the coating solution may be from about 1% to about 20% by weight of the dry polymer.

The method of coating the granular core particles comprises applying aqueous dispersions of Eudragit RL30D and RS30D to the core as distinct layers in subsequent coating steps. The polymers will be present in the dispersions in an amount of from about 10% to about 30% as the dry polymer. Advantageously, the core material is coated by subjecting them to air suspension coating technique. Preferably a Glatt Fluid Bed Wurster Coater is employed. An inner coating is applied by spraying (bottom spray) onto the fluidizing drug cores an aqueous dispersion of RL30D. The spraying is continued until a theoretical weight gain of from about 25% to about 50% of the batch is obtained. The coated material is retrieved and dedusted on a #60-80 U.S. mesh screen, equivalent to approximately 180-250 microns.

A second coating layer comprising an aqueous dispersion of RS30D is applied to the above coated, dedusted particles to a theoretical weight gain of from about 5% to about 15% of the original batch. When the coating run is completed the particles are again dedusted using a #80 mesh screen resulting in a final product having a particle size of from about 200 to about 400 microns.

The dedusting between coating steps is preferable particularly when friable drug particles are employed. More important, the dedusting steps result in a drug particle size which contributes to the immediate release and bioavailability characteristics of the coated drug granules.

One of the advantages of the coated drug particles of this invention is that they provide a method for formulating extremely palatable solid dosage units which contain

unpleasant tasting drugs. For example, orally administrable dosage units such as chewable tablets, troches, lozenges or sprinkle formulations may be prepared from the coated granules of this invention. The granules of the present invention are particularly suitable for use in preparing chewable tablets.

- 5           Where the dosage unit is a chewable tablet, conventional pharmaceutical excipients in addition to the coated core may be included. For example, fillers, lubricants, binders, compression aids, and wetting agents may be employed. The fillers may be water soluble or insoluble such as lactose, sucrose or terra alba. Typical lubricants are stearic acid and its pharmaceutically acceptable alkali metal salts.
- 10       Examples of binders are polyvinylpyrrolidones, polyethylene glycol, natural gums including veegum, tragacanth, acacia and gelatin. Exemplary of wetting agents include sodium lauryl sulfate, polysorbates and polyoxyethylene surfactants. Starch, alginates and their salts and maize would be representative of disintegrants. Typical compression aids would be microcrystalline cellulose, dicalcium phosphate and compressible sugar.
- 15       To further assist patient compliance, the tablet can also contain sweeteners such as aspartame, sodium cyclamate and sodium saccharinate and flavorants such as orange, mint, cola, strawberry and the like.

The invention is illustrated but in no way limited by the following examples.

20

Example 1

Eudragit RL30D - Inner Coating Solution

<u>Ingredients</u>	<u>Amount/Gms</u>
Eudragit RL30D (dry polymer)	160.00
Triethyl Citrate	32.00
25   Talc	1.60
Yellow #6 Lake	1.60
Distilled Water	768.00

- 30           To 533.33 gms of a 30% aqueous dispersion of Eudragit RL30D (equivalent to 160.00 gms of dry polymer) was added the triethyl citrate with gentle mixing. The talc and lake were then added with low shear mixing. The distilled water was added and slowly mixed with the suspension.

Eudragit RS30D Outer Coating Solution

	<u>Ingredients</u>	<u>Amount/Gms.</u>
	Eudragit RS30D (dry polymer)	40.00
5	Triethyl citrate	8.00
	Yellow #6 Lake	0.40
	Talc	0.40
	Distilled Water	192.00

- 10 To 133.33 gms of a 30% aqueous dispersion of RS30D (equivalent to 40.00 gms of dry polymer) the ingredients were added and mixed following the above procedure.

Preparation of Coated Granules

- 15 Employing a Glatt Fluid Bed Wurster Coater, 400 mg of caffeine having a size of from about 180 to 420 microns were fluidized. Onto the fluidizing caffeine a first coating of the above Eudragit RL30D aqueous dispersion was applied by the bottom spray until a 40% weight gain of the batch was obtained. The batch was retrieved and dedusted on a #60 mesh screen to remove the fines. The sifted coated particles were then returned to the Wurster chamber and a second coating of the above RS30D
- 20 aqueous dispersion was applied by the bottom spray until a 10% weight gain of the original batch was achieved. The coated particles were then removed, dedusted on a #60 mesh screen resulting in a final particle size of 200 to 400 microns.

Preparation of Chewable Tablet

25	<u>Ingredients</u>	<u>Amount/mg</u>
	Caffeine coated beads*	400.00
	Compressible Sugar N.F.	1595.00
	Magnesium Stearate	500.00
	*Equivalent to 200 mg. of Caffeine	

30

The ingredients were thoroughly mixed and directly compressed employing the appropriate punches and dies.



Example 2Eudragit RL 30D - Inner Coating Solution

	<u>Ingredients</u>	<u>Amount/Gms.</u>
5	Eudragit RL 30D (dry polymer)	248.00
	Triethyl citrate	49.60
	Talc	2.48
	Distilled Water	1121.78

- 10 To 826.67 g of a 30% aqueous dispersion of Eudragit RL 30D (equivalent to 248.00 g of dry polymer) was added the triethyl citrate with gentle mixing. The talc was then added with low shear mixing. The distilled water was added and slowly mixed with the suspension.

15 Eudragit RS 30D - Outer Coating Solution

	<u>Ingredients</u>	<u>Amount/Gms.</u>
	Eudragit RS 30D (dry polymer)	62.00
	Triethyl citrate	12.40
	Yellow #6 Lake	0.62
20	Talc	0.62
	Distilled Water	283.96

- 25 To 206.67 g of a 30% aqueous dispersion of RS 30D (equivalent to 62.00 g of dry polymer) was added the triethyl citrate with gentle mixing. The talc and lake were then added with low shear mixing. The distilled water was added and slowly mixed with the suspension.

Preparation of Coated Granules

- 30 Employing a Glatt Fluid Bed Wurster Coater, 600.0 gms of cimetidine having a size of from about 250 to 177 microns were fluidized. Onto the fluidized cimetidine a first coating of the above Eudragit RL 30D aqueous dispersion was applied by the bottom spray to obtain a target weight gain of 40%. The batch was retrieved and sieved on a #60 mesh screen to remove the fines. The sifted coated particles were then returned to the Wurster chamber and a second coating of the above RS 30D aqueous

dispersion was applied by the bottom spray to obtain a target weight gain of 10% of the original batch. The coating particles were then removed, dedusted on a # 60 mesh screen resulting in a final particle size of about 200 to 400 microns.

5	<u>Preparation of Chewable Tablet</u>	
	<u>Ingredients</u>	<u>Amount in Milligrams</u>
	Cimetidine coated granules *	161.0
	Compressible Sugar N.F.	1829.0
	Magnesium Stearate	10.0

10

\* Equivalent to 100.0 mg of Cimetidine

The ingredients were thoroughly mixed and directly compressed employing the appropriate punches and dies.

15

Unexpectedly, it has been discovered that when the polymers RL30D and RS30D are applied as distinct coating layers as described hereinabove they not only produce extremely palatable granules which significantly minimizes the unpleasant taste associated with bitter drugs but also result in an immediate release of the drug in the stomach. This is surprising because both polymers are employed to produce either a

20

sustained release or an enteric coated effect of the drug.

A further advantage of this invention is that by employing an aqueous based coating medium the dangers associated with organic solvents, such as, for example, flammability, toxicity and pollution of the environment are eliminated.

The coating system of this invention releases the drug by diffusion and not

25

explosion. The release is influenced by drug solubility and media pH.

What is claimed is:

- 5           1. A taste masked pharmaceutical granule composition for oral  
administration comprising a core material containing an unpleasant tasting drug  
said core having an inner polymeric coating of high permeability comprising  
poly(ethyl acrylate-methyl methacrylate) trimethylammonio ethyl methacrylate  
chloride in a 1:2:0.2 ratio and a outer polymeric coating of low permeability  
10 comprising poly(ethyl acrylate-methyl methacrylate) trimethylammonio ethyl  
methacrylate chloride in a 1:2:0.1 ratio said coating providing an immediate release  
of the drug in the stomach.
- 15           2. The composition of Claim 1 wherein said polymeric coating contains a  
plasticizer.
3. The composition of Claim 2 wherein the plasticizer is present in an  
amount of from about 5% to about 30% and the coated granules have a final  
particle size of from about 200 to about 400 microns.  
20
4. The composition of Claims 1, 2 or 3 wherein said drug is selected from  
the group consisting of cimetidine, ranitidine, caffeine, acetaminophen,  
dextromethorphan, chlorpheniramine, diphenhydramine, theophylline, penicillin,  
ampicillin and erythromycin.  
25
5. A chewable tablet containing the pharmaceutical granule composition of  
Claim 1 and optionally further contains a pharmaceutical excipient.
6. A chewable tablet containing the granule composition of Claim 4.  
30
7. The chewable tablet of Claim 6 wherein the drug is cimetidine and it is  
present in an amount of from about 100 mg to about 400 mg.
8. A method for preparing a taste masked pharmaceutical granule  
35 composition which comprises spraying onto a fluidized bed of a drug core material  
an aqueous dispersion of an inner polymeric coating of high permeability,  
retrieving and dedusting the coated core material and applying to a fluidized bed of

the dedusted material an aqueous dispersion of an outer polymeric coating of low permeability dedusting and drying the coated particles.

5

9. The method of Claim 8 wherein the inner polymeric coating is poly(ethylacrylate-methyl methacrylate) trimethylammonio ethyl methacrylate chloride in a 1:2:0.2 ratio and the polymeric outer coating is poly(ethylacrylate methylmethacrylate)trimethylammonio ethyl melthacrylate chloride in a 1:2:0.1 ratio and the coated material is dedusted to a final particle size of from about 200 to about 400 microns.

10

10. The method of Claim 8 wherein the drug is cimetidine or caffeine.

# INTERNATIONAL SEARCH REPORT

Int. l. application No.  
PCT/US93/05423

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 9/16  
US CL :424/497

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/497, 472, 470

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,075,114 (ROCHE) 24 DECEMBER 1991 See entire document.	1-4, 7-10
Y	US, A, 5,112,621 (STEVENS) 12 MAY 1992 See col. 1, lines 20-41, 57-59.	5-6



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be part of particular relevance	* X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* E* earlier document published on or after the international filing date	* Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* &*	document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means		
* P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

21 JULY 1993

Date of mailing of the international search report

NOV 12 1993

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

WILLIAM E. BENSTON, JR.

Telephone No. (703) 308-2351

**THIS PAGE BLANK (USPTO)**